

LETTERS AND  
CORRESPONDENCE

*Letters and correspondence submitted for possible publication must be identified as such. Text length must not exceed 500 words and five bibliographic references. A single concise figure or table may be included if it is essential to support the communication. Letters not typed double-spaced will not be considered for publication. Letters not meeting these specifications will not be returned to authors. Letters to the Editor are utilized to communicate a single novel observation or finding. Correspondence is to be used to supplement or constructively comment on the contents of a publication in the journal and cannot exceed the restrictions for Letters to the Editor. The Editor reserves the right to shorten text, delete objectional comments, and make other changes to comply with the style of the journal. Permission for publication must be appended as a postscript. Submissions must be sent to Paul Chervenick, M.D., Editor of Brief Reports/Letters to Editors, American Journal of Hematology, H. Lee Moffitt Cancer Center, University of South Florida, 12902 Magnolia Drive, Tampa, FL 33612 to permit rapid consideration for publication.*

### Arterial Thrombosis, GM-CSF, and the Lupus Anticoagulant

*To the Editor:* Recombinant colony stimulating factors such as granulocyte-macrophage colony stimulating factor (GM-CSF) have been advocated for use in severe drug-induced neutropenia. Side effects include both arterial and venous thrombosis, although almost all reported cases have occurred in cancer patients on chemotherapy. We report a case of arterial thrombosis in a patient with clozapine-induced neutropenia being treated with GM-CSF, who had the potentially additive prothrombotic risk factor of a drug-induced lupus anticoagulant (LAC).

A 30-year-old polynesian male with longstanding manic depression was admitted 8 weeks after starting clozapine, his neutrophil count having fallen to  $0.1 \times 10^9/L$ . He had been treated in previous years with fluphenazine and chlorpromazine. He was a smoker with no family history of thrombosis. Treatment with oral ciprofloxacin and fluconazole antimicrobial prophylaxis, and GM-CSF (300  $\mu g/day$  subcutaneously) was started. On day 3 of treatment he suddenly developed a dense left-sided hemiplegia. CT scan showed a large cerebral infarction in the right middle cerebral artery territory. Doppler studies showed complete occlusion of right internal and external carotid arteries with thrombus. Activated partial thromboplastin time was prolonged at 39 sec (control 36 sec), and prothrombin time was normal. LAC testing (by Kaolin Clotting Time and Dilute Russell's Viper Venom Test) was positive. Screening for anticardiolipin antibodies, protein S, protein C, and antithrombin III deficiencies, as well as

activated protein C resistance was negative. Fibrinogen was initially elevated at 8.5 g/L (later falling to 6.0 g/L). ANA was detectable at a titre of 1:40; double-stranded DNA was negative.

GM-CSF was replaced with G-CSF, and the patient was warfarinized.

### DISCUSSION

There have been many case reports of thrombotic events in cancer patients on chemotherapy receiving GM-CSF for neutropenia. A recent meta-analysis strongly suggested (but did not prove) that GM-CSF, but less so granulocyte colony stimulating factor (G-CSF), was associated with thrombotic risk [1]. There are no reports of thrombotic events in drug-induced neutropenia treated with GM-CSF that we are aware of. Potential mechanisms of thrombosis include increased tissue factor expression on macrophages [2], enhanced expression of adhesion molecules on neutrophils (with resultant binding to vascular endothelium resulting in damage) [1], and increased platelet aggregation (reported with G-CSF) [3].

Given that the majority of reported cases had additional thrombotic risk factors it is of interest that our patient was also discovered to have a (presumed) drug-induced lupus anticoagulant (LAC). Chlorpromazine has been associated with the LAC in up to 50% of treated patients and a case of clozapine-induced LAC was recently reported in this journal [4]. The risk of thrombosis in these patients is controversial, and while some authors are convinced that it is increased [5] the overall incidence appears to be low. It may, however, have contributed to our patient's catastrophic thrombotic event.

Given that G-CSF appears to have a lesser prothrombotic tendency than GM-CSF, we would advocate the use of the former where potential thrombotic risk factors already exist.

**MARK. W. DRUMMOND**  
**RUTH SPEARING**

*Department of Haematology, Western Infirmary, Glasgow, Scotland, UK*

### REFERENCES

1. Barbui T, Finazzi G, Grassi A, Marchioli R. Thrombosis in cancer patients treated with haematopoietic growth factors: a meta analysis. On behalf of the Subcommittee on Haemostasis and Malignancy of the Scientific and Standardisation Committee of the ISTH. *Thrombosis Haemostasis* 1996;75(2):368–371.
2. Edwards RL, Rickles FR. The role of leucocytes in the activation of blood coagulation. *Semin Haematol* 1992;29:202–212.
3. Kuriowa M, Okanura T, Kanaji T, Okamura S, Harada M, Niho Y. Effects of granulocyte colony-stimulating factor on the haemostatic system in healthy volunteers. *Int J Hematol* 1996;63(4):311–316.
4. Kanjolia A, Valigorsky JM, Josion-Pasion ML. Clozaril-induced lupus anticoagulant. *Am J Hematol* 1997;54:345–346.
5. Bick RL, Baker WF. Antiphospholipid and thrombosis syndromes. *Semin Thrombosis Haemostasis* 1994;20:3–15.